

Study of temporal quantum correlations in decohering B and K meson systems

Javid Naikoo,^{*} Ashutosh Kumar Alok,[†] and Subhashish Banerjee[‡]
Indian Institute of Technology Jodhpur, Jodhpur 342011, India

(Dated: February 14, 2018)

In this work we study temporal quantum correlations, quantified by Leggett-Garg (LG) and LG-type inequalities, in the B and K meson systems. We use the tools of open quantum systems to incorporate the effect of decoherence which is quantified by a single phenomenological parameter. The effect of CP violation is also included in our analysis. We find that the LG inequality is violated for both B and K meson systems, the violation being most prominent in the case of K mesons and least for B_s system. Since the systems with no coherence do not violate LGI, incorporating *decoherence* is expected to decrease the extent of violation of LGI and is clearly brought out in our results. We show that the expression for the LG functions depends upon an additional term, apart from the experimentally measurable meson transition probabilities. This term vanishes in the limit of zero decoherence. On the other hand, the LG-type parameter can be directly expressed in terms of transition probabilities, making it a more appropriate observable for studying temporal quantum correlations in neutral meson systems.

I. INTRODUCTION

Quantum correlations (QCs), existing between two or more parties [1, 2], are bestowed with properties unique to the quantum world and are of pivotal importance in quantum information science. The study of QCs not only unveils the fundamental traits responsible for the distinction of the quantum mechanically correlated systems from those attributed with a joint classical probability distribution [3], it also helps in devising efficient ways of carrying out the tasks of quantum communication and computation [4–6].

Among the most celebrated notions in quantum physics are nonlocality [7], entanglement [1], quantum discord [8, 9], and teleportation fidelity [10, 11]. These spatial quantum correlations (SQC) have enhanced our understanding of nature at the fundamental level and at the same time have provided efficacious solutions in the development of the theory of quantum information. The SQCs mentioned above have been studied in many systems, viz., optical systems [12–18], NMR [19–21], neutrino oscillation [22–25], B and K meson systems [26]. Of the above listed SQCs, Bell nonlocality is the strongest and Bell inequalities are considered to be the oldest tool for detecting entanglement [27].

The temporal quantum correlations (TQCs) arising from the sequential measurements on a system at different times, have also been considered as promising candidates in discerning the quantum behavior from the classical. Leggett and Garg inequalities (LGIs) [28] are among the well known TQCs, violation of which is a witness of quantum *coherence* in the system. LGIs have been a topic of study in various theoretical works [29–35] including, in recent times, neutrino oscillations [25, 36, 37] and studied experimentally in systems like superconducting qubits [38, 39], photons [40–43], and NMR [44–46].

Leggett-Garg inequalities are based on the concept of *macrorealism* (MR) and *noninvasive measurability* (NIM). MR means that the system which has available to it two or more macroscopically distinct states, pertaining to an observable \hat{Q} , always exists in one of these states irrespective of any measurement performed on it. NIM states that, in principle, we can perform the measurement without disturbing the future dynamics of the system [34]. MR and NIM put limits on certain combinations of the two time correlation functions $C_{ij} = \langle Q(\hat{t}_i)Q(\hat{t}_j) \rangle$. Quantum systems, however, violate these limits. The simplest form of LGI is the one involving three measurements performed at t_0 , t_1 and t_2 ($t_2 > t_1 > t_0$)

$$K_3 = C_{01} + C_{12} - C_{02}, \quad (1)$$

such that $-3 \leq K_3 \leq 1$. The maximum quantum value of K_3 for a two level system is $\frac{3}{2}$ [28] and has been found to hold for any system, *irrespective* of the number of levels, as long as the measurements are given by just two projectors Π^\pm [47], a fact revealed in several studies [32, 48–50]. It was shown in [51] that in the limit $N \rightarrow \infty$, the LGI can be violated up to its maximum algebraic sum.

^{*}Electronic address: naikoo.1@iitj.ac.in

[†]Electronic address: akalok@iitj.ac.in

[‡]Electronic address: subhashish@iitj.ac.in

The autocorrelation C_{12} turns out to contain a nonmeasurable quantity and hence reduces the efficacy of Eq. (1) from the experimental point of view. Such limitations of the two time correlations have been discussed in detail in [52–55], and a different approach was developed which involves replacing the NIM by a weaker condition called *stationarity*. This avoids the need of performing the measurement at the intermediate time t_1 by replacing C_{12} by C_{01} , thus leading to an easily testable Leggett-Garg type inequality (LGtI)

$$\tilde{K}_3 = 2C_{01} - C_{02} \leq 1. \quad (2)$$

The following set of assumptions [34] are considered important for applying *stationarity* to a system: (i) macroscopic realism, (ii) the conditional probability $P(\psi, t + t_0 | \psi, t_0)$ of finding the system in state ψ at time $t + t_0$ given that it was in state ψ at time t_0 , should be invariant under the time translation, $P(\psi, t + t_0 | \psi, t_0) = P(\psi, t | \psi, 0)$, (iii) Markovianity and (iv) that the system is prepared in state ψ at time $t = 0$.

In this work, we study the LG and LG-type inequalities in the B and K meson systems. The effect of decoherence is included by using the formalism of open quantum systems. Decoherence, here, is modelled by a single phenomenological parameter [56] which represents the interaction between the one-particle system and its environment. The environment can be attributed to quantum gravity effects [57–64] or it can be due to detector background itself. Apart from decoherence, we also include the effects of CP violation. We find that the LG inequality is violated for both B and K meson systems. Apart from the experimentally measurable meson transition probabilities, we show that the LG function depends upon an additional term which vanishes in the limit of zero decoherence. The LG-type parameter on the other hand can be directly expressed in terms of transition probabilities.

The plan of this work is as follows. In the next section, we discuss the time evolution of B and K meson systems treated as open quantum systems. In Sec. III, we derive the LG and LG-type inequalities for these systems. In Sec. IV, we present our results. Finally, in Sec. V, we make our conclusions.

II. B AND K MESONS AS OPEN QUANTUM SYSTEMS

In this section, we introduce our formalism for the study of B and K mesons as open quantum systems.

A. Kraus representation

Kraus representation [65], describes the time evolution of an *open* quantum system, which is not necessarily unitary unlike the evolution of a *closed* quantum system. Real physical systems are always entangled with their ambient environment, alternately addressed as the reservoir. Kraus representations are very convenient for handling a number of practical problems of open system dynamics [66–71]. Consider a large system S comprising of two subsystems S_a and S_b . At a given time t , let the quantum states corresponding to S , S_a and S_b be represented by $\rho(t)$, $\rho_a(t)$ and $\rho_b(t)$, respectively. Then $\rho_a(t) = Tr_b\{\rho(t)\}$ and $\rho_b(t) = Tr_a\{\rho(t)\}$. Since the total system is unitary, its evolution is given by

$$\rho(t) = U(t)\rho(0)U^\dagger(t), \quad (3)$$

where $U(t)$ is a unitary operator. The evolution of system S_a will look like

$$\rho_a(t) = Tr_b\{U(t)\rho(0)U^\dagger(t)\}. \quad (4)$$

If it is possible to recast Eq. (4) in the following form

$$\rho_a(t) = \sum_i E_i(t)\rho_a(0)E_i^\dagger(t), \quad (5)$$

such that $\sum_i E_i(t)E_i^\dagger(t) = \mathbb{1}$, then the evolution of $\rho_a(t)$ has a *Kraus* representation and is completely positive.

B. Time evolution of B/K mesons

We describe briefly the time evolution of $B^0(K^0)$ meson system. Since both B^0 and K^0 share the same scheme of dynamics, we discuss only B^0 system and the results, with appropriate notational changes, will be applicable to the K^0 system. The states of the total system, including the meson and the vacuum $|0\rangle$, introduced in order to

incorporate the effect of decay in the meson system, reside in the Hilbert space given by the direct sum $\mathcal{H}_{B^0} \oplus \mathcal{H}_0$ [56, 72, 73] spanned by the orthonormal vectors $|B^0\rangle$, $|\bar{B}^0\rangle$ and $|0\rangle$

$$|B^0\rangle = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}; \quad |\bar{B}^0\rangle = \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}; \quad |0\rangle = \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}. \quad (6)$$

Here B^0 stands for B_d^0/B_s^0 mesons. The mass eigenstates $\{|B_L\rangle, |B_H\rangle\}$ are related to the flavor eigenstates $\{|B^0\rangle, |\bar{B}^0\rangle\}$ by the equations

$$|B_L\rangle = p|B^0\rangle + q|\bar{B}^0\rangle, \quad |B_H\rangle = p|B^0\rangle - q|\bar{B}^0\rangle, \quad (7)$$

with $|p|^2 + |q|^2 = 1$. The time evolution is given by a family of completely positive trace preserving maps forming a one parameter dynamical semigroup. The complete positivity requires the time evolution of a state of the system being represented by the operator-sum representation [65]

$$\rho(t) = \sum_{i=0} E_i(t)\rho(0)E_i^\dagger(t), \quad (8)$$

where the *Kraus* operators have the following form

$$\begin{aligned} E_0 &= |0\rangle\langle 0|, \\ E_1 &= \mathcal{E}_{1+}(|B^0\rangle\langle B^0| + |\bar{B}^0\rangle\langle \bar{B}^0|) + \mathcal{E}_{1-}\left(\frac{p}{q}|B^0\rangle\langle \bar{B}^0| + \frac{q}{p}|\bar{B}^0\rangle\langle B^0|\right), \\ E_2 &= \mathcal{E}_2\left(\frac{p+q}{2p}|0\rangle\langle B^0| + \frac{p+q}{2q}|0\rangle\langle \bar{B}^0|\right), \\ E_3 &= \mathcal{E}_{3+}\frac{p+q}{2p}|0\rangle\langle B^0| + \mathcal{E}_{3-}\frac{p+q}{2q}|0\rangle\langle \bar{B}^0|, \\ E_4 &= \mathcal{E}_4(|B^0\rangle\langle B^0| + |\bar{B}^0\rangle\langle \bar{B}^0| + \frac{p}{q}|B^0\rangle\langle \bar{B}^0| + \frac{q}{p}|\bar{B}^0\rangle\langle B^0|), \\ E_5 &= \mathcal{E}_5(|B^0\rangle\langle B^0| + |\bar{B}^0\rangle\langle \bar{B}^0| - \frac{p}{q}|B^0\rangle\langle \bar{B}^0| - \frac{q}{p}|\bar{B}^0\rangle\langle B^0|). \end{aligned}$$

Here the coefficients are

$$\mathcal{E}_{1\pm} = \frac{1}{2} \left[e^{-(2im_L + \Gamma_L + \lambda)t/2} \pm e^{-(2im_H + \Gamma_H + \lambda)t/2} \right], \quad (9a)$$

$$\mathcal{E}_2 = \sqrt{\frac{Re[\frac{p-q}{p+q}]}{|p|^2 - |q|^2} (1 - e^{-\Gamma_L t} - (|p|^2 - |q|^2)^2 \frac{1 - e^{-(\Gamma + \lambda - i\Delta m)t}}{1 - e^{-\Gamma_H t}})}, \quad (9b)$$

$$\mathcal{E}_{3\pm} = \sqrt{\frac{Re[\frac{p-q}{p+q}]}{(|p|^2 - |q|^2)(1 - e^{-\Gamma_H t})}} [1 - e^{-\Gamma_H t} \pm (1 - e^{-(\Gamma + \lambda - i\Delta m)t})(|p|^2 - |q|^2)], \quad (9c)$$

$$\mathcal{E}_4 = \frac{e^{-\Gamma_L t/2}}{2} \sqrt{1 - e^{-\lambda t}}, \quad (9d)$$

$$\mathcal{E}_5 = \frac{e^{-\Gamma_H t/2}}{2} \sqrt{1 - e^{-\lambda t}}. \quad (9e)$$

A meson initially in state $\rho_{B^0}(0) = |B^0\rangle\langle B^0|$ or $\rho_{\bar{B}^0}(0) = |\bar{B}^0\rangle\langle \bar{B}^0|$, after time t , evolves to

$$\rho_{B^0}(t) = \frac{1}{2} e^{-\Gamma t} \begin{pmatrix} a_{ch} + e^{-\lambda t} a_c & \left(\frac{q}{p}\right)^* (-a_{sh} - ie^{-\lambda t} a_s) & 0 \\ \left(\frac{q}{p}\right) (-a_{sh} + ie^{-\lambda t} a_s) & |q/p|^2 a_{ch} - e^{-\lambda t} a_c & 0 \\ 0 & 0 & \rho_{33}(t) \end{pmatrix}, \quad (10)$$

and

$$\rho_{B^0}(t) = \frac{1}{2}e^{-\Gamma t} \begin{pmatrix} \frac{|p|}{q}|^2(a_{ch} - e^{-\lambda t}a_c) & (\frac{p}{q})(-a_{sh} + ie^{-\lambda t}a_s) & 0 \\ (\frac{p}{q})^*(-a_{sh} - ie^{-\lambda t}a_s) & a_{ch} + e^{-\lambda t}a_c & 0 \\ 0 & 0 & \tilde{\rho}_{33}(t) \end{pmatrix}. \quad (11)$$

Here, a_{ch} (a_{sh}) and a_c (a_s) stand for the hyperbolic functions $\cosh[\frac{\Delta\Gamma t}{2}]$ ($\sinh[\frac{\Delta\Gamma t}{2}]$) and the trigonometric functions $\cos[\Delta mt]$ ($\sin[\Delta mt]$), respectively. p and q are defined in Eq. (7). $\Delta\Gamma = \Gamma_L - \Gamma_H$ is the difference of the decay width Γ_L (for B_L^0) and Γ_H (for B_H^0). $\Gamma = \frac{1}{2}(\Gamma_L + \Gamma_H)$ is the average decay width. The mass difference $\Delta m = m_H - m_L$, where m_H and m_L are the masses of B_H^0 and B_L^0 states, respectively. The strength of the interaction between the one particle system and its environment is quantified by λ , the *decoherence* parameter [74]. The elements $\rho_{33}(t)$ and $\tilde{\rho}_{33}(t)$ are known functions of B physics parameters, not used in this work. In the following section, we use this formalism to develop the LGI and LGtI for the meson systems.

III. TEMPORAL QUANTUM CORRELATIONS IN B/K SYSTEMS

A. Leggett-Garg inequality

Leggett-Garg inequalities, often referred to as the temporal Bell inequalities, place bounds on certain combinations of the two time autocorrelations C_{ij} , defined in terms of the joint probabilities as [28, 75, 76]

$$C_{ij} = p(+t_i)q(+t_j|+t_i) - p(+t_i)q(-t_j|+t_i) - p(-t_i)q(+t_j|-t_i) + p(-t_i)q(-t_j|-t_i), \quad (12)$$

where $p(+t_i)$ is the probability of obtaining the result $a = \pm 1$ at t_i , and $q(+t_j|+t_i)$ is the conditional probability of getting result $b = \pm 1$ at time t_j , given that result $a = \pm 1$ was obtained at t_i . To find the probabilities involved in Eq. (12), we define the projector Π^\pm related to the eigenspace of the dichotomic operator \hat{Q} , such that the probability of obtaining outcome a at time t_i is

$$p(+t_i) = Tr\{\Pi^+ \rho(t_i)\} = Tr\{\Pi^+ \sum_{\mu} K_{\mu}(t_i) \rho(0) K_{\mu}^{\dagger}(t_i)\}. \quad (13)$$

The density matrix corresponding to the measurement result a obtained at ' t_i ' is given by the von Neumann rule

$$\rho^a(t_i) = \frac{\Pi^a \rho(t_i) \Pi^a}{Tr\{\Pi^a \rho(t_i)\}} = \frac{\Pi^a \sum_{\mu} K_{\mu}(t_i) \rho(0) K_{\mu}^{\dagger}(t_i) \Pi^a}{p(+t_i)}, \quad (14)$$

this state evolves until t_j , when the state of the system looks like $\sum_{\nu} K_{\nu}(t_j - t_i) \rho^a(t_i) K_{\nu}^{\dagger}(t_j - t_i)$, so that the probability of obtaining outcome b at time t_j , given that a was obtained at time t_i , is given by

$$q(+t_j|+t_i) = \frac{Tr\{\Pi^+ \sum_{\nu, \mu} K_{\nu}(t_j - t_i) \Pi^+ K_{\mu}(t_i) \rho(0) K_{\mu}^{\dagger}(t_i) \Pi^+ K_{\nu}^{\dagger}(t_j - t_i)\}}{p(+t_i)}. \quad (15)$$

A generic term in the right-hand side of Eq. (12) becomes

$$p(+t_i)q(+t_j|+t_i) = Tr\{\Pi^+ \sum_{\nu} K_{\nu, \mu}(t_j - t_i) \Pi^+ K_{\mu}(t_i) \rho(0) K_{\mu}^{\dagger}(t_i) \Pi^+ K_{\nu}^{\dagger}(t_j - t_i)\}. \quad (16)$$

With some algebra, we can show that the two time correlations turn out to be [76]

$$C_{ij} = 1 - 2p(+t_1) - 2p(+t_2) + 4Re[g(t_i, t_j)], \quad (17)$$

where

$$g(t_i, t_j) = Tr\left\{\Pi^+ \sum_{\nu} K_{\nu}(t_j - t_i) \Pi^+ \rho(t_i) K_{\nu}^{\dagger}(t_j - t_i)\right\}. \quad (18)$$

We consider a dichotomic quantity $Q = \pm 1$ for our *three* level system, such that each level is associated with a definite value of Q . Assigning the same value of Q to different states is irrelevant from the macrorealistic point

of view and does not change the bounds of Eq. (1) [51]. Let us assume that at time $t = 0$, the meson was in state $\rho_{\bar{B}^0}$. This state evolves to $\rho_{\bar{B}^0(t_i)}$ at time t_i and is given by Eq. (11). We define the dichotomic operator $\Pi = \Pi^+ - \Pi^- = \Pi_{B^0} - (\Pi_{\bar{B}^0} + \Pi_0)$, where $\Pi_x = |x\rangle\langle x|$. Now

$$p(^+t_i) = Tr\{\Pi^+ \rho_{\bar{B}^0}(t_i)\} = [\rho_{\bar{B}^0}(t_i)]_{11} = |p/q|^2 \frac{e^{-\Gamma t_i}}{2} \left[\cosh\left(\frac{\Delta\Gamma t_i}{2}\right) - e^{-\lambda t_i} \cos(\Delta m t_i) \right]. \quad (19)$$

Thus, $p(^+t_i) = \mathcal{P}_{\bar{B}^0 B^0}(t_i)$ is the transition probability from state $\rho_{\bar{B}^0}$ to ρ_{B^0} at time t_i . With the assumption of equal time measurements $t_2 - t_1 = t_1 - 0 = \Delta t$, we have the following expression for C_{12}

$$C_{12} = 1 - 4\mathcal{P}_{\bar{B}^0 B^0}(\Delta t) + 4Re[g(\Delta t)], \quad (20)$$

with

$$g(t_1, t_2) = 2\mathcal{P}_{\bar{B}^0 B^0}(\Delta t)\mathcal{P}_{\bar{B}^0 B^0}(\Delta t) + \left|\frac{p}{q}\right|^2 \frac{e^{-2\Gamma\Delta t}(e^{-2\lambda\Delta t} - 1)}{4}. \quad (21)$$

Here $\mathcal{P}_{\bar{B}^0 \bar{B}^0}(\Delta t)$ and $\mathcal{P}_{\bar{B}^0 B^0}(\Delta t)$ are the survival and transition probabilities, respectively, for the meson which started in state $\rho_{\bar{B}^0} = |\bar{B}^0\rangle\langle\bar{B}^0|$ at time $t = 0$. The survival probability of \bar{B}^0 has the following form:

$$\mathcal{P}_{\bar{B}^0 \bar{B}^0}(t) = \frac{e^{-\Gamma t}}{2} \left[\cosh\left(\frac{\Delta\Gamma t}{2}\right) + e^{-\lambda t} \cos(\Delta m t) \right]. \quad (22)$$

The LG function finally becomes

$$K_3 = 1 - 4\mathcal{P}_{\bar{B}^0 B^0}(\Delta t) + 8\mathcal{P}_{\bar{B}^0 B^0}(\Delta t)\mathcal{P}_{\bar{B}^0 \bar{B}^0}(\Delta t) + |p/q|^2 e^{-2\Gamma\Delta t}(e^{-2\lambda\Delta t} - 1). \quad (23)$$

CP violation implies that $|p/q| \neq 1$. The above developed formalism also applies to the K meson case with some notational changes. The CP violating parameter for K mesons ϵ can be expressed in terms of p and q by the following relation $\epsilon = \frac{p-q}{p+q}$.

B. Leggett-Garg type inequality

The assumption of noninvasive measurability makes it difficult to test the Leggett-Garg inequality experimentally. Different measurement strategies like negative outcome measurement, delayed choice measurement, weak measurements [38, 40, 77–79] have been devoted to this effect. Another formalism developed in [52, 53], replacing the assumption of noninvasive measurability by “stationarity”, leads to easily testable inequalities using projective (von Neumann) measurements. According to the stationarity assumption, the conditional probability $q(t_i, t_j)$ to find a system in state j at time t_j , if it was in state i at time t_i only depends on the time difference $t_j - t_i$, and is expected to hold not only for idealized closed quantum systems, but also in open quantum systems subjected to purely Markovian noise at a rate γ such that the two time correlations are exponentially damped by a factor $\gamma(t_2 - t_1)$ [55]. The full set of assumptions (i)-(iv), for the stationarity to hold for a system, as given in Sec. (I), turns out to be applicable in the context of K and B meson systems. Given that the state of the meson at time $t = 0$ is $|\bar{B}^0\rangle$, it can be shown that Markovian dynamics described by the Kraus operators in Sec. (II) lead to the time translation invariance of the conditional probability, i.e., $P(\bar{B}^0, t + t_0 | \bar{B}^0, t_0) = P(\bar{B}^0, t | \bar{B}^0, 0)$. With the assumption of stationarity, the Leggett-Garg type inequality, Eq. (2), becomes

$$\tilde{K}_3 = 1 - 4\mathcal{P}_{\bar{B}^0 B^0}(\Delta t) + 2\mathcal{P}_{\bar{B}^0 B^0}(2\Delta t). \quad (24)$$

Therefore, a knowledge of the transition probabilities at times Δt and $2\Delta t$ would allow one to compute K_3 according to Eq. (24), such that $\tilde{K}_3 > 1$ shows the nonclassical nature of the neutral meson oscillations. It should be noted that Eq. (24) is expressed completely in terms of directly measurable quantities such as transition probabilities unlike Eq. (23), which contains a term $(|p/q|^2 e^{-2\Gamma\Delta t}(e^{-2\lambda\Delta t} - 1))$, apart from the survival and transition probabilities. However, it can be seen that in the limit of neglecting decoherence effects, Eq. (23), can also be expressed directly in terms of survival and transition probabilities.

The experiments on the $B^0(K^0)$ meson systems involve determination of their flavor at the time of production or decay. This is done by analyzing the flavor specific decays. For e.g., a B_d^0 meson can decay into a positron (or a μ^+), a neutrino and a hadron with a branching ratio of ~ 0.1 . This semileptonic decay is induced by the quark level transition $\bar{b} \rightarrow \bar{c} l^+ \nu_l$, with $l = e, \mu$. On the other hand, the corresponding decay of a \bar{B}_d^0 meson results in an electron




FIG. 1: The left, middle and right panels of the figure depict the LG function K_3 plotted *w.r.t* the dimensionless quantity $\Delta t/\tau$ for the K , B_d and B_s mesons, respectively. Here Δt is the time between successive measurements and τ is the mean lifetime of respective mesons. Dashed and solid curves correspond to the cases with and without decoherence, respectively. For the K system, the mean lifetime is $\tau_K = 1.7889 \times 10^{-10} s$. Also, $\Gamma = 5.59 \times 10^9 s^{-1}$, $\Delta\Gamma = 1.1174 \times 10^{10} s^{-1}$, $\lambda = 2.0 \times 10^8 s^{-1}$ and $\Delta m = 5.302 \times 10^9 s^{-1}$ [80]. Here we used $Re(\epsilon) = 1.596 \times 10^{-3}$ and $|\epsilon| = 2.228 \times 10^{-3}$ [81]. For the B_d system, $\tau_{B_d} = 1.518 \times 10^{-12} s$, $\Gamma = 6.58 \times 10^{11} s^{-1}$, $\Delta\Gamma = 0$, $\lambda = 0.012 \times 10^{12} s^{-1}$ and $\Delta m = 0.5064 \times 10^{12} s^{-1}$ [82]. The CP violating parameter used here is $|\frac{q}{p}| = 1.010$ [82]. Finally, for the B_s meson, $\tau_{B_s} = 1.509 \times 10^{-12} s$, $\Gamma = 0.6645 \times 10^{12} s^{-1}$, $\Delta\Gamma = 0.086 \times 10^{12} s^{-1}$, $\lambda = 0.012 \times 10^{12} s^{-1}$ and $\Delta m = 17.757 \times 10^{12} s^{-1}$ [82]. The value of the CP violating parameter here is $\frac{q}{p} = 1.003$ [82]. As we do not have any experimental bound on the decoherence parameter λ for the B_s system, we assume it to be the same as that of the B_d system.

(or a μ^-) in the final state. Thus, in general, the charge of the final state lepton is same as the charge of the decaying quark. This is known as the $\Delta B = \Delta Q$ rule for the semileptonic decays of B mesons and is assumed in most of the experimental analysis. Hence, the charge of the final state lepton in the semi-leptonic decays of a neutral meson usually determines the flavor of that meson at the time of decay.

The process of determination of the initial flavor of a neutral meson is called tagging. This is achieved by making use of the rule of associated production. The mesons are produced either by strong or electromagnetic interactions and hence a quark is always produced in association with its anti-quark as flavor is conserved in these interactions. Thus, if a quark q is detected at one end of the detector then at the quark at the other end has to be \bar{q} . Now if a charged meson is produced in association with a neutral meson, then the decay of the charged meson determines the flavor of the neutral meson at production. This is so because the charged meson cannot oscillate. The survival and oscillation probability of the neutral meson can then be measured by identifying the charge of the lepton in its semileptonic decay. If two entangled neutral mesons are produced, as in the e^+e^- colliders by the process $e^+e^- \rightarrow \Upsilon(4S) \rightarrow B_d^0 \bar{B}_d^0$, then detecting the flavor specific final state of one meson, say at time t_1 , determines the flavor of that meson as well as the other meson at that time t_1 . The oscillation probability of the tagged meson is then determined by identifying its final flavor specific state.

IV. RESULTS AND DISCUSSION

The left panel of Fig. (1) shows the variation of the LG function K_3 , as a function of the dimensionless quantity $\Delta t/\tau_K$. It can be seen from the figure that the LG inequality is violated for about $\Delta t = \tau_K$. The middle and right panels of Fig. (1) depict the variation of the LG function for the B_d and B_s mesons, respectively. One can see that the violation in the B_d meson system sustains for about $\Delta t = \tau_{B_d}$ while for the B_s meson system the violation is roughly for $\Delta t \approx 0.5 \tau_{B_s}$. The maximum violation of LGI occurs around $\Delta t \approx 0.41\tau_K$, $\Delta t \approx 0.37\tau_{B_d}$ and $\Delta t \approx 0.037\tau_{B_s}$ for K , B_d and B_s meson systems, respectively.

The figures clearly bring out the point that from the genesis of its decay [73], the meson systems violate the upper threshold value of $K_3 = 1$, indicative of quantum behavior, and quickly fall below one. The K_3 value for K meson remains above one longest while B_s does it for the shortest time. In addition, the B_s meson exhibits an additional recurrence behavior. In order to have an understanding of this recurrence behavior, we re-write Eq. (23) as

$$K_3 = 1 + |p/q|^2 \left[2e^{-(\Gamma+\lambda)\Delta t} \cos(\Delta m \Delta t) - e^{-2(\Gamma+\lambda)\Delta t} \cos(2\Delta m \Delta t) - 2e^{-\Gamma\Delta t} \cosh(\Delta\Gamma\Delta t/2) + e^{-2\Gamma\Delta t} \cosh(\Delta\Gamma\Delta t) \right]. \quad (25)$$

One can then see that the oscillating behavior in the case of B_s meson system could be attributed to the mass term Δm (Eq. (25)), which plays the role of frequency, and is more than 35 times the corresponding value for the B_d

meson system.

From Eq. (24), we find that the LG-type inequality is in terms of the transition probabilities only. Fig. (2) shows the deviation of the LG-type function, \tilde{K}_3 (24), from the LG-function (K_3). It is clear from the figure that the deviation is very small. Thus, a study of the LG inequality in mesons, using \tilde{K}_3 , Eq. (24), in terms of experimentally measurable quantities would be well justified. Eq. (24) demands the knowledge of the transition probabilities at Δt and $2\Delta t$, for example, $(0.5\tau_K, \tau_K)$ for the K meson system.




FIG. 2: Plot of the difference of LG-function K_3 and LG-type function \tilde{K}_3 in the case of K meson system. The various parameters used are the same as in Fig. (1).

Looking at the form of Eq. (23), it can be seen that the only nonmeasurable term in the equation is $|p/q|^2 e^{-2\Gamma\Delta t} (e^{-2\lambda\Delta t} - 1)$; we call this term \mathcal{D}_B and \mathcal{D}_K for the case of B meson and K meson systems, respectively. In the limit of zero decoherence, $\lambda \rightarrow 0$, $\mathcal{D}_{B/K} \rightarrow 0$, rendering the LG function, Eq. (23), in terms of measurable survival and transition probabilities

$$K_3(\lambda = 0) = 1 - 4P_{B^0 B^0}(\Delta t) + 8P_{B^0 B^0}(\Delta t)P_{\bar{B}^0 \bar{B}^0}(\Delta t). \quad (26)$$

The variation of \mathcal{D}_B and \mathcal{D}_K with $\Delta t/\tau_{K/B_{d(s)}}$ is shown in Fig. 3. It is obvious from the figure that these terms




FIG. 3: The nonmeasurable term $\mathcal{D}_K = |(1 + \epsilon)/(1 - \epsilon)|^2 e^{-2\Gamma\Delta t} (e^{-2\lambda\Delta t} - 1)$ for K -meson system and $\mathcal{D}_{B_{d(s)}} = |p/q|^2 e^{-2\Gamma\Delta t} (e^{-2\lambda\Delta t} - 1)$ for $B_{d(s)}$ -meson system, plotted against $\Delta t/\tau_{K/B_{d(s)}}$. The various parameters used in the two cases are the same as mentioned in the caption of Fig. (1).

are small compared with the maximum value attained by the LG function K_3 .

V. CONCLUSION

In this work, we study the violation of LG and LG-type inequalities in B and K mesons within the framework of open quantum systems. It is found that LGI is violated in both K and B meson systems. This violation lasts for a longer time in the case of K mesons as compared to that of B mesons. In the case of B meson systems, the violation lasts longer for B_d mesons as compared to the B_s system. We show that the LG function K_3 , apart from the measurable survival and transition probabilities, contains a nonmeasurable term which is small compared to the maximum value attained by it and vanishes in the approximation of zero decoherence. Since systems with no coherence do not violate LGI, the effect of *decoherence* should result in decreasing the extent of the violation, as observed in Fig. (1). Further, it is highlighted in this work that the LG-type function, unlike LG function, can be

expressed completely in terms of experimentally measurable quantities. Hence, LG type inequality is seen to be more suitable for understanding the nature of temporal quantum correlations in meson systems.

-
- [1] R. Horodecki, P. Horodecki, M. Horodecki, and K. Horodecki, *Rev. Mod. Phys.* **81**, 865 (2009).
- [2] K. Modi, A. Brodutch, H. Cable, T. Paterek, and V. Vedral, *Rev. Mod. Phys.* **84**, 1655 (2012).
- [3] J. F. Clauser and A. Shimony, *Rep. Prog. Phys.* **41**, 1881 (1978).
- [4] N. Gisin, G. Ribordy, W. Tittel, and H. Zbinden, *Rev. Mod. Phys.* **74**, 145 (2002).
- [5] R. Raussendorf and H. J. Briegel, *Phys. Rev. Lett.* **86**, 5188 (2001).
- [6] H. J. Briegel, D. E. Browne, W. Dür, R. Raussendorf, and M. Van den Nest, *Nat. Phys.* **5**, 19 (2009).
- [7] J. S. Bell, *Physics (Long Island City, N.Y.)* **1**, 195 (1964).
- [8] H. Ollivier and W. H. Zurek, *Phys. Rev. Lett.* **88**, 017901 (2001).
- [9] L. Henderson and V. Vedral, *J. Phys. A* **34**, 6899 (2001).
- [10] M. Barrett, J. Chiaverini, T. Schaetz, J. Britton, W. Itano, J. Jost, E. Knill, C. Langer, D. Leibfried, R. Ozeri, *et al.*, *Nature (London)* **429**, 737 (2004).
- [11] S. Adhikari and S. Banerjee, *Phys. Rev. A* **86**, 062313 (2012).
- [12] A. Aspect, P. Grangier, and G. Roger, *Phys. Rev. Lett.* **47**, 460 (1981).
- [13] W. Tittel, J. Brendel, B. Gisin, T. Herzog, H. Zbinden, and N. Gisin, *Phys. Rev. A* **57**, 3229 (1998).
- [14] W. Tittel, J. Brendel, H. Zbinden, and N. Gisin, *Phys. Rev. Lett.* **81**, 3563 (1998).
- [15] G. Weihs, T. Jennewein, C. Simon, H. Weinfurter, and A. Zeilinger, *Physical Review Letters* **81**, 5039 (1998).
- [16] I. Chakrabarty, S. Banerjee, and N. Siddharth, *Quantum Information and Computation* **11**, 0541 (2011).
- [17] B. Lanyon, P. Jurcevic, C. Hempel, M. Gessner, V. Vedral, R. Blatt, and C. Roos, *Phys. Rev. Lett.* **111**, 100504 (2013).
- [18] J. Naikoo, K. Thapliyal, A. Pathak, and S. Banerjee, arXiv:1712.04154 ().
- [19] A. R. Kessel and V. L. Ermakov, arXiv preprint quant-ph/0011002 (2000).
- [20] K. Dorai, A. Kumar, *et al.*, *Pramana* **56**, L705 (2001).
- [21] R. Laflamme, D. G. Cory, C. Negrevergne, and L. Viola, quant-ph/0110029 .
- [22] M. Blason, F. Dell'Anno, S. De Siena, and F. Illuminati, *Europhys. Lett.* **85**, 50002 (2009).
- [23] A. K. Alok, S. Banerjee, and S. U. Sankar, *Nucl. Phys. B* **909**, 65 (2016).
- [24] S. Banerjee, A. K. Alok, R. Srikanth, and B. C. Hiesmayr, *Eur. Phys. J. C* **75**, 487 (2015).
- [25] J. Naikoo, A. K. Alok, S. Banerjee, S. U. Sankar, G. Guarneri, and B. C. Hiesmayr, arXiv:1710.05562 ().
- [26] S. Banerjee, A. K. Alok, and R. MacKenzie, *Eur. Phys. J. Plus* **131**, 1 (2016).
- [27] O. Gühne and G. Tóth, *Phys. Rep.* **474**, 1 (2009).
- [28] A. J. Leggett and A. Garg, *Phys. Rev. Lett.* **54**, 857 (1985).
- [29] M. Barbieri, *Phys. Rev. A* **80**, 034102 (2009).
- [30] D. Avis, P. Hayden, and M. M. Wilde, *Phys. Rev. A* **82**, 030102 (2010).
- [31] N. Lambert, C. Emary, Y.-N. Chen, and F. Nori, *Phys. Rev. Lett.* **105**, 176801 (2010).
- [32] N. Lambert, R. Johansson, and F. Nori, *Phys. Rev. B* **84**, 245421 (2011).
- [33] A. Montina, *Phys. Rev. Lett.* **108**, 160501 (2012).
- [34] C. Emary, N. Lambert, and F. Nori, *Rep. Prog. in Phys.* **77**, 016001 (2013).
- [35] J. Kofler and Č. Brukner, *Phys. Rev. A* **87**, 052115 (2013).
- [36] J. Formaggio, D. Kaiser, M. Murskyj, and T. Weiss, *Phys. Rev. Lett.* **117**, 050402 (2016).
- [37] Q. Fu and X. Chen, *Eur. Phys. J. C* **77**, 775 (2017).
- [38] A. Palacios-Laloy, F. Mallet, F. Nguyen, P. Bertet, D. Vion, D. Esteve, and A. N. Korotkov, *Nat. Phys.* **6**, 442 (2010).
- [39] J. Groen, D. Ristè, L. Tornberg, J. Cramer, P. C. De Groot, T. Picot, G. Johansson, and L. DiCarlo, *Phys. Rev. Lett.* **111**, 090506 (2013).
- [40] M. Goggin, M. Almeida, M. Barbieri, B. Lanyon, J. O'Brien, A. White, and G. Pryde, *Proc. Nat. Acad. Sci.* **108**, 1256 (2011).
- [41] J.-S. Xu, C.-F. Li, X.-B. Zou, and G.-C. Guo, *Scientific Reports* **1** (2011).
- [42] J. Dressel, C. Broadbent, J. Howell, and A. N. Jordan, *Phys. Rev. Lett.* **106**, 040402 (2011).
- [43] Y. Suzuki, M. Inuma, and H. F. Hofmann, *New J. Phys.* **14**, 103022 (2012).
- [44] V. Athalye, S. S. Roy, and T. Mahesh, *Phys. Rev. Lett.* **107**, 130402 (2011).
- [45] A. Souza, I. Oliveira, and R. Sarthour, *New J. Phys.* **13**, 053023 (2011).
- [46] H. Katiyar, A. Shukla, K. R. K. Rao, and T. Mahesh, *Phys. Rev. A* **87**, 052102 (2013).
- [47] C. Budroni, T. Moroder, M. Kleinmann, and O. Gühne, *Phys. Rev. Lett.* **111**, 020403 (2013).
- [48] R. E. George, L. M. Robledo, O. J. Maroney, M. S. Blok, H. Bernien, M. L. Markham, D. J. Twitchen, J. J. Morton, G. A. D. Briggs, and R. Hanson, *Proc. Nat. Acad. Sci.* **110**, 3777 (2013).
- [49] J. Kofler and Č. Brukner, *Phys. Rev. Lett.* **99**, 180403 (2007).
- [50] M. M. Wilde, J. M. McCracken, and A. Mizel, *Pro. R. Soc. A* **466**, 1347 (2010).
- [51] C. Budroni and C. Emary, *Phys. Rev. Lett.* **113**, 050401 (2014).
- [52] S. F. Huelga, T. W. Marshall, and E. Santos, *Physical Review A* **52**, R2497 (1995).
- [53] S. F. Huelga, T. W. Marshall, and E. Santos, *Physical Review A* **54**, 1798 (1996).

- [54] S. Huelga, T. Marshall, and E. Santos, *Europhysics. Lett.* **38**, 249 (1997).
- [55] G. Waldherr, P. Neumann, S. Huelga, F. Jelezko, and J. Wrachtrup, *Phys. Rev. Lett.* **107**, 090401 (2011).
- [56] A. K. Alok, S. Banerjee, and S. Uma Sankar, *Phys. Lett. B* **749**, 94 (2015).
- [57] S. W. Hawking, *Comm. Math. Phys.* **87**, 395 (1982).
- [58] J. R. Ellis, J. S. Hagelin, D. V. Nanopoulos, and M. Srednicki, *Nucl. Phys.* **B241**, 381 (1984).
- [59] J. R. Ellis, N. Mavromatos, and D. V. Nanopoulos, *Int. J Mod. Phys. A* **13**, 1059 (1998).
- [60] N. E. Mavromatos and S. Sarkar, *Phys. Rev. D* **74**, 036007 (2006).
- [61] X.-K. Guo, arXiv:1512.05277 .
- [62] J. c. v. Minář, P. Sekatski, and N. Sangouard, *Phys. Rev. A* **94**, 062111 (2016).
- [63] B. O. Kerbikov, arXiv:1704.07117 .
- [64] A. Roberts, *Phys. Rev. D* **96** (2017).
- [65] K. Kraus, *States, Effects and Operations* (Springer, New York, 1983).
- [66] H.-P. Breuer and F. Petruccione, *The Theory of Open Quantum Systems* (Oxford University Press on Demand, New York, 2002).
- [67] U. Weiss, *Quantum Dissipative Systems*, Vol. 13 (World Scientific, Singapore, 2012).
- [68] M. A. Nielsen and I. Chuang, *Quantum Computation and Quantum Information*. Cambridge University Press, Cambridge, England (2000).
- [69] S. Banerjee, V. Ravishankar, and R. Srikanth, *Eur. Phys. J. D* **56**, 277 (2010).
- [70] S. Banerjee, V. Ravishankar, and R. Srikanth, *Ann. Phys. (N.Y.)* **325**, 816 (2010).
- [71] R. S. S. Omkar and S. Banerjee, *Quantum Information Processing* **12**, 3725 (2013).
- [72] P. Caban, J. Rembieliński, K. A. Smoliński, and Z. Walczak, *Phys. Rev. A* **72**, 032106 (2005).
- [73] A. K. Alok and S. Banerjee, *Phys. Rev. D* **88**, 094012 (2013).
- [74] A. K. Alok, S. Banerjee, and S. Uma Sankar, *Phys. Lett. B* **749**, 94 (2015).
- [75] J. Kofler and Č. Brukner, *Phys. Rev. Lett.* **101**, 090403 (2008).
- [76] J. C. Castillo, F. J. Rodríguez, and L. Quiroga, *Phys. Rev. A* **88**, 022104 (2013).
- [77] C. Tesche, *Phys. Rev. Lett.* **64**, 2358 (1990).
- [78] J. P. Paz and G. Mahler, *Phys. Rev. Lett.* **71**, 3235 (1993).
- [79] A. Fedrizzi, M. P. Almeida, M. A. Broome, A. G. White, and M. Barbieri, *Phys. Rev. Lett.* **106**, 200402 (2011).
- [80] C. Patrignani *et al.*, *Chin. Phys C* **40**, 100001 (2016).
- [81] G. D'Ambrosio and G. e. a. K. c. Isidori, *J. High Energy Phys.* **2006**, 011 (2006).
- [82] Y. Amhis *et al.*, *Eur. Phys. J. C* **77**, 895 (2017).