

Laplacian matrices of weighted digraphs represented as quantum states*

Bibhas Adhikari[†] Subhashish Banerjee[‡] Satyabrata Adhikari[§]
and Atul Kumar[¶]

Abstract

Graph representation of quantum states is becoming an increasingly important area of research to investigate combinatorial properties of quantum states which are nontrivial to comprehend in standard linear algebraic density matrix based approach of quantum states. In this paper, we propose a general weighted directed graph framework for investigating properties of a large class of quantum states which are defined by three types of Laplacian matrices associated with such graphs. We generalize the standard framework of defining density matrices from simple connected graphs to density matrices using both combinatorial and signless Laplacian matrices associated with weighted directed graphs with complex edge weights and with/without loops. We also introduce a new notion of Laplacian matrix which we call signed Laplacian matrix associated with such graphs. We produce necessary and/or sufficient conditions for such graphs to represent pure and mixed quantum states. Using these criteria we finally determine the graphs whose corresponding density matrices represent entangled pure states which are well-known and important for quantum computation applications. It is important to observe that all these entangled pure states share a common combinatorial structure.

Keywords: Combinatorial Laplacian, signless Laplacian, eigenvalues, pure and mixed states, density matrix, quantum entanglement

1 Introduction

Quantum mechanics deals with states living in the Hilbert space, allowing for linear superpositions to be built up, a facility of immense importance for harnessing the power of quantum mechanics but at the same time making it computationally a formidable task. This can be most easily appreciated by considering entanglement [1, 2] in higher dimensions as well as in multi-partite systems [3], all mathematically and computationally very formidable tasks. Any tool that would aid in this regard would be very welcome.

In combinatorics a graph is a collection of vertices and edges which link two vertices. A digraph is a graph consisting directed edges. A loop is an edge that joins a vertex with itself. A graph is said to be a weighted graph if each edge is assigned a nonzero number which is called the weight of the corresponding edge. The theory of graphs is a well-developed mathematical theory that has found many applications in diverse areas, such as the spectrum of a discrete Schrödinger operator in a uniform, periodic magnetic field [4]. Graphs have, by their very construction, the inherent feature of visualization. A pertinent question to ask is whether graphical representation of quantum states can be made? This would enable

*This work is supported by CSIR (Council of Scientific and Industrial Research) Grant No. 25(0210)/13/EMR-II, New Delhi, India.

[†]Department of Mathematics, IIT Kharagpur, India, E-mail: bibhas@maths.iitkgp.ernet.in

[‡]Department of Physics, IIT Jodhpur, India, E-mail: subhashish@iitj.ac.in

[§]Department of Mathematics, BIT Mesra, India, E-mail: tapisatya@gmail.com

[¶]Department of Chemistry, IIT Jodhpur, India, E-mail: atulk@iitj.ac.in

the incorporation of the mathematical machinery of graphs into the problems of quantum mechanics and at the same time bring in the attractive feature of visualization of quantum states.

Attempts have been made to realize density matrix representation of a quantum state by defining a matrix associated with a graph. In this case, the graph is called the graph representation of a quantum state. This idea was first introduced in [5] by considering a combinatorial Laplacian matrix associated with an unweighted undirected graph (simple graph). It was further extended in [6] for weighted graphs. A criteria of separability of multipartite states represented by the combinatorial Laplacian matrices of simple graphs have also developed in [7]. Recently, local unitary transformations on a density matrix obtained by signless Laplacian matrix associated with a simple graph has been established as a combinatorial operation which is known as switching of a graph in [8]. A combinatorial operation has also been introduced for density matrices defined by Laplacian matrices associated with simple graphs in [9] that act as an entanglement generator for mixed states arising from partially symmetric graphs.

In this paper, we use both combinatorial and signless Laplacian matrices to define density matrices associated with a weighted digraph having complex edge weights and with or without loops. We also introduce a matrix, which we call signed Laplacian matrix associated with a weighted digraph having loops with both positive and negative weights. In order to relate the topological structure of a weighted digraph and properties of the density matrices defined by these Laplacian matrices, we investigate the zero eigenvalues of these matrices associated with such graphs. This leads to a classification of graphs which always provide density matrices representing pure quantum states, and those which determine mixed quantum states. Indeed, since the number of mixed states are significantly larger than the number of pure states, emphasis is given on identifying graphs which can produce important classes of pure states. We also provide graphs which identify a number of well-known entangled pure states by using the Laplacian matrices associated with such graphs. A state vector of an entangled state cannot be expressed as a tensor product of other state vectors. These states play an important role in different tasks of quantum information theory. We observe that in the graph representation of entangled pure states, all the weighted edges are clustered in a subgraph which forms a completely connected graph and the weight of the loops attached at each of the vertices is $-(m-2)$ where m is the number of vertices involved in the complete subgraph. This should be of interest to the quantum information community.

The plan of the paper is as follows. In section 2, we define the required terminologies of graph theory and in particular, we investigate the Laplacian matrices and Laplacian spectra associated with a weighted digraph with or without loops. In section 3, we define the density matrix associated with a weighted digraph and classify the graphs which represent pure and mixed states. Finally, we provide graphs which define density matrices of entangled pure states.

2 Weighted digraphs with/without loops and its Laplacian spectra

Let $G = (V, E)$ be a graph with the vertex set $V = \{1, 2, \dots, n\}$ and edge set $E \subseteq V \times V$. A directed graph or digraph G is a graph with a function assigning to each edge an ordered pair of vertices. The first vertex of the ordered pair is called the initial vertex of the edge, and the second is the terminal vertex; together, they are the endpoints. Thus each edge of a digraph is directed; and an undirected edge can be considered as both way directed. A weighted graph G is a graph with a function $w : E \rightarrow \mathbb{C}$, defined by $w(i, j) = w_{ij}$, $1 \leq i, j \leq n$, where \mathbb{C} is the set of all complex numbers. If $|w_{ij}| = 1$ for all $(i, j) \in E$, such a graph is called gain graph [10]. The function w is called an edge-weight function and w_{ij} is called the weight of $(i, j) \in E$. An unweighted graph can be considered as an edge-weighted graph with weight

function $w_{ij} = 1$ if $w_{ij} \neq 0$. A weighted digraph is a graph which is both weighted and directed. However, as mentioned, an unweighted undirected graph also can be considered as weighted digraph with constant weight function with weight of any edge is 1 and the edges are both way directed. Thus, from now onward a graph always is a weighted digraph unless otherwise mentioned.

2.1 Weighted digraphs with or without loops having nonnegative weights

First, we consider weighted digraphs having loops (at least one loop and maximum one loop at a vertex) with nonnegative real weights. The adjacency matrix $A(G) = (a_{ij})$ associated with G is defined as

$$a_{ij} = \begin{cases} w_{ij}, & \text{if } (i, j) \in E; \\ \overline{w_{ij}}, & \text{if } (j, i) \in E; \\ r_i, & \text{if } (i, i) \in E; \\ 0, & \text{otherwise} \end{cases}$$

where $0 \leq r_i \in \mathbb{R}$ is the weight of the loop at i th vertex and $w_{ij} \in \mathbb{C}$. Note that $r_i = 0$ if i th vertex does not have a loop. The weighted degree d_i of a vertex $i \in V$ is given by $d_i = \sum_{j=1}^n |a_{ij}|$. The Laplacian and the signless Laplacian matrices are defined by

$$L(G) = \text{diag}(\{d_i\}_{i=1}^n) - A(G) \text{ and } Q(G) = \text{diag}(\{d_i\}_{i=1}^n) + A(G), \quad (1)$$

respectively [11, 12, 13]. Notice that loops, even though apparent in the adjacency matrix $A(G)$, do not appear in the Laplacian matrix $L(G)$. The above constructions of $L(G)$ and $Q(G)$ will lead to diagonally dominant matrices, i.e., a matrix M where $|M_{i,i}| \geq \sum_{j \neq i} |M_{i,j}|$.

For a weighted edge from vertex i to j with weight $w_{ij} \in \mathbb{C}$, we assume $w_{ij} = r_{ij} e^{i\theta_{ij}}$, $r_{ij} > 0$, $0 \leq \theta \leq \pi$. We consider $\overline{w_{ij}} = r_{ij} e^{-i\theta_{ij}}$ - $\pi \leq \theta_{ij} \leq 0$, $\sqrt{w_{ij}} = \sqrt{r_{ij}} e^{i\theta_{ij}/2}$, $\sqrt{\overline{w_{ij}}} = \sqrt{r_{ij}} e^{-i\theta_{ij}/2}$. Thus, $(\sqrt{\overline{w_{ij}}})^2 = w_{ij}$ and $\sqrt{w_{ij}} \sqrt{\overline{w_{ij}}} = r_{ij}$.

Lemma 2.1 *Let $G = (V, E)$ be a weighted directed graph without loops. Then $L(G)$ and $Q(G)$ are Hermitian and positive semidefinite matrices.*

Proof: Assume that $w_{ij} \in \mathbb{C}$ is the weight of an edge (i, j) . Then define

$$(M^-)_{v,e} = \begin{cases} \sqrt{w_{ij}}, & \text{if } w_{ij} \in \mathbb{C} \setminus \mathbb{R} \text{ and } v \text{ is initial vertex of nonloop edge } e \\ -\sqrt{\overline{w_{ij}}}, & \text{if } w_{ij} \in \mathbb{C} \setminus \mathbb{R} \text{ and } v \text{ is terminal vertex of nonloop edge } e \\ \sqrt{|w_{ij}|}, & \text{if } 0 > w_{ij} \in \mathbb{R} \text{ and } v \text{ is initial or terminal vertex of nonloop edge } e \\ -\sqrt{|w_{ij}|}, & \text{if } 0 < w_{ij} \in \mathbb{R} \text{ and } v \text{ is initial vertex of nonloop edge } e \\ \sqrt{|w_{ij}|}, & \text{if } 0 < w_{ij} \in \mathbb{R} \text{ and } v \text{ is terminal vertex of nonloop edge } e \\ 0, & \text{otherwise} \end{cases}$$

and

$$(M^+)_{v,e} = \begin{cases} \sqrt{w_{ij}}, & \text{if } w_{ij} \in \mathbb{C} \setminus \mathbb{R} \text{ and } v \text{ is initial vertex of nonloop edge } e \\ \sqrt{\overline{w_{ij}}}, & \text{if } w_{ij} \in \mathbb{C} \setminus \mathbb{R} \text{ and } v \text{ is terminal vertex of nonloop edge } e \\ \sqrt{|w_{ij}|}, & \text{if } 0 < w_{ij} \in \mathbb{R} \text{ and } v \text{ is initial or terminal vertex of nonloop edge } e \\ -\sqrt{|w_{ij}|}, & \text{if } 0 > w_{ij} \in \mathbb{R} \text{ and } v \text{ is initial vertex of nonloop edge } e \\ \sqrt{|w_{ij}|}, & \text{if } 0 > w_{ij} \in \mathbb{R} \text{ and } v \text{ is terminal vertex of nonloop edge } e \\ 0, & \text{otherwise} \end{cases}$$

for any $v \in V$ and $e \in E$. Then it is easy to verify that $L(G) = M^-(M^-)^\dagger$ and $Q(G) = M^+(M^+)^\dagger$ where † denotes conjugate transpose. Therefore, the result follows.

From the construction of M^+ and M^- in Lemma 2.1, it follows that

$$\begin{aligned} x^\dagger L(G)x = & \sum_{i \neq j, (i,j) \in E, 0 < w_{ij} \in \mathbb{R}} |w_{ij}| |x_i - x_j|^2 + \sum_{i \neq j, (i,j) \in E, 0 > w_{ij} \in \mathbb{R}} |w_{ij}| |x_i + x_j|^2 \\ & + \sum_{i \neq j, (i,j) \in E, w_{ij} \in \mathbb{C} \setminus \mathbb{R}} |\sqrt{w_{ij}} x_i - \sqrt{w_{ij}} x_j|^2 \end{aligned} \quad (2)$$

and

$$\begin{aligned} x^\dagger Q(G)x = & \sum_{i \neq j, (i,j) \in E, 0 < w_{ij} \in \mathbb{R}} |w_{ij}| |x_i + x_j|^2 + \sum_{i \neq j, (i,j) \in E, 0 > w_{ij} \in \mathbb{R}} |w_{ij}| |x_i - x_j|^2 \\ & + \sum_{i \neq j, (i,j) \in E, w_{ij} \in \mathbb{C} \setminus \mathbb{R}} |\sqrt{w_{ij}} x_i + \sqrt{w_{ij}} x_j|^2 \end{aligned} \quad (3)$$

In the following theorem we provide a necessary and sufficient condition for a connected loopless weighted digraph having a signless Laplacian eigenvalue zero. Recall that, a digraph is said to be connected if it is connected without considering the directions of the edges.

Theorem 2.2 *Let G be a connected weighted digraph without selfloops. The least eigenvalue of the signless Laplacian matrix of a loopless connected weighted digraph is 0 if and only if*

$$(-1)^{p_+ + |P|} \prod_{(i,j) \in P, w_{ij} \in \mathbb{C} \setminus \mathbb{R}} \frac{w_{ij}}{|w_{ij}|} = (-1)^{p'_+ + |P'|} \prod_{(i,j) \in P', w_{ij} \in \mathbb{C} \setminus \mathbb{R}} \frac{w_{ij}}{|w_{ij}|}$$

holds for any two directed paths P and P' with numbers of links having positive real weights are p_+ and p'_+ respectively between any fixed two vertices and $|P|$ denotes the number of edges having nonreal weights in P .

Proof: Assume that the least eigenvalue of the signless Laplacian $Q(G)$ of G has an eigenvalue zero, that is, $x^H Q(G)x = 0$ for some non-zero vector x . From (3), it is obvious that for such x , $x_i = -x_j$ if $0 < w_{ij} \in \mathbb{R}$; $x_i = x_j$ if $0 > w_{ij} \in \mathbb{R}$; $x_i = -\frac{w_{ij}}{|w_{ij}|} x_j$ if $w_{ij} \in \mathbb{C} \setminus \mathbb{R}$. Let $P \equiv (u = i_1, i_2, \dots, i_{k_1} = v)$ and $P' \equiv (u = i'_1, i'_2, i'_3, \dots, i'_{k_2} = v)$ be two distinct directed paths from the vertex u to the vertex v . Then, for the path P ,

$$x_u = (-1)^{p_+ + |P|} \prod_{(i,j) \in P, w_{ij} \in \mathbb{C} \setminus \mathbb{R}} \frac{w_{ij}}{|w_{ij}|} x_v;$$

and for the path P' ,

$$x_u = (-1)^{p'_+ + |P'|} \prod_{(i,j) \in P', w_{ij} \in \mathbb{C} \setminus \mathbb{R}} \frac{w_{ij}}{|w_{ij}|} x_v.$$

Further, $x_u \neq 0$ and $x_v \neq 0$ since otherwise $x = 0$ follows from (3) as the graph is connected. Hence the desired result follows.

Conversely, if the given condition is true for any two different directed paths for any pair of vertices in G , a vector x defined by $x_i = -x_j$ if $0 < w_{ij} \in \mathbb{R}$; $x_i = x_j$ if $0 > w_{ij} \in \mathbb{R}$; $x_i = -\frac{w_{ij}}{|w_{ij}|} x_j$ if $w_{ij} \in \mathbb{C} \setminus \mathbb{R}$ will satisfy $x^\dagger Q(G)x = 0$. Hence the proof.

Corollary 2.3 *The least eigenvalue of the signless Laplacian of a loopless connected weighted digraph having complex unit weights is equal to 0 if and only if*

$$W(P) = (-1)^{p' - p} W(P')$$

holds for any two directed walks P, P' of lengths p and p' respectively between any fixed two vertices where $W(P)$ (resp. $W(P')$) is the product of the weights of the edges of P (resp. P'). In particular, 0 is a simple eigenvalue.

Corollary 2.4 *Let G be a weighted digraph without loops with $n(> 2)$ vertices. Assume that 0 is a signless Laplacian eigenvalue of G . Then the multiplicity of 0 as a signless Laplacian eigenvalue of G is k if and only if the graph is disconnected with k connected components.*

It is shown in [13] that for a unweighted undirected connected graph G without loops, the least eigenvalue of the signless Laplacian of G is equal to 0 if and only if the graph is bipartite and 0 is a simple eigenvalue. We mention that the condition provided in Theorem 2.3 for existence of zero eigenvalue of weighted directed graph is a generalized version of the condition obtained for unweighted undirected graph in [13]. That is, the condition in Theorem 2.3 is satisfied for an unweighted undirected connected graph if and only if the graph is bipartite.

The following corollary provides a necessary and sufficient condition for existence of zero Laplacian eigenvalue of a weighted connected digraph.

Corollary 2.5 *The least eigenvalue of the combinatorial Laplacian matrix of a loopless connected weighted digraph is equal to 0 if and only if*

$$(-1)^{p_-} \prod_{(i,j) \in P, w_{ij} \in \mathbb{C} \setminus \mathbb{R}} \frac{w_{ij}}{|w_{ij}|} = (-1)^{p'_-} \prod_{(ij) \in P', w_{ij} \in \mathbb{C} \setminus \mathbb{R}} \frac{w_{ij}}{|w_{ij}|}$$

holds for any two directed walks P, P' with number of links with negative real weights arcs p_- and p'_- respectively between any fixed two vertices.

Proof: The proof is similar to the proof of Theorem 2.2.

Corollary 2.6 *The least eigenvalue of the combinatorial Laplacian of a loopless connected weighted digraph having complex unit weights is equal to 0 if and only if*

$$W(P) = W(P')$$

holds for any two walks P, P' between any two vertices where $W(P)$ (resp. $W(P')$) is the product of the weights of the edges of P (resp. P'). In particular, 0 is a simple eigenvalue.

An alternative proof of the above corollary also can be found in [12].

Corollary 2.7 *Let G be a weighted digraph without loops with $n(> 2)$ vertices. Assume that 0 is a combinatorial Laplacian eigenvalue of G . Then the multiplicity of 0 as a combinatorial Laplacian eigenvalue of G is k if and only if the graph is disconnected with k connected components.*

As we mentioned above, loops with nonnegative weights have no effect on the combinatorial Laplacian matrix. Thus, we only consider signless Laplacian matrix when an weighted digraph contains at least one loop with positive real weight. It is easy to verify that, given an weighted digraph G with at least one loop having positive real weight, we have

$$x^\dagger Q(G)x = x^\dagger Q(\widehat{G})x + \sum_{(i,i) \in E} r_i |x_i|^2 \quad (4)$$

where \widehat{G} is the subgraph of G without considering loops. This also shows that $Q(G)$ is Hermitian and positive semidefinite.

Lemma 2.8 *0 can never be a signless Laplacian eigenvalue of a connected weighted digraph G with loops (at least one vertex contains a loop) having positive weights.*

Proof: Consider a connected weighted digraph G with loops (at least one vertex contains a loop) having positive weights. If 0 is a signless Laplacian eigenvalue of G , from (4) we know that, there exists an $0 \neq x \in \mathbb{C}^n$ such that

$$x^\dagger Q(G)x = x^\dagger Q(\widehat{G})x + \sum_{(i,i) \in E} r_i |x_i|^2 = 0$$

where \widehat{G} is the subgraph of G without loops. Assume that the k -th vertex contains the loop. For $x^\dagger Q_G x$ to be zero, x_k has to be zero since r_k is positive. Further, since the graph is connected, k th vertex is linked with m (say) vertices k_1, k_2, \dots, k_m for some m which implies $x_{k_j} = 0$ for $j = 1, \dots, m$ which further implies that $x_j = 0$ for all $j = 1, \dots, n$ since k_j vertices are linked with other vertices and $x^\dagger Q(\widehat{G})x = 0$ for all $(i, j) \in E$.

2.2 Weighted digraphs with loops having at least one loop with negative weight

Recall the definitions of Laplacian and signless Laplacian matrices associated with a weighted digraph. Observe that loops, even though apparent in the adjacency matrix $A(G)$ do not reflect in Laplacian matrix when the loops are having positive weights, and in signless Laplacian matrices when the loops are having negative weights. Thus, for weighted digraphs with both nonnegative and negative weighted loops (at least one of the loops has negative weight), we introduce a new matrix, which we call *signed Laplacian*, denoted by $L_-(G)$ and $L_\pm(G)$ when G has all the loops with nonpositive weights, and when G contains loops with positive weights as well as negative weights, respectively by

$$L_-(G) = R_-(G) + Q(\widehat{G}) \text{ and } L_\pm(G) = R_\pm(G) + Q(\widehat{G}) \quad (5)$$

where $R_-(G) = R_\pm(G) = \text{diag}\{r_1, r_2, \dots, r_n\}$, $r_j \in \mathbb{R}$ denotes the weight of the loop at j th vertex, $j = 1, \dots, n$ and $Q(\widehat{G})$ is the signless Laplacian matrix of the graph \widehat{G} constructed from G without considering the loops. Obviously, $L_-(G)$ and $L_\pm(G)$ are Hermitian matrices. In order to simplify the notation we denote $R(G) = R_-(G) = R_\pm(G)$.

Lemma 2.9 *Given a weighted digraph G with nonpositive loops, $L_-(G)$ is semi-definite if*

$$\max_i |r_i| \leq \lambda_{\min} Q(\widehat{G})$$

where $r_i \leq 0, i = 1 : n$ are the weights of the loops present in the graph and $\lambda_{\min} Q(\widehat{G})$ denotes the minimum eigenvalue of $Q(\widehat{G})$.

Proof: For any unit vector $0 \neq x \in \mathbb{C}^n$, we have

$$-x^\dagger R(G)x \leq \max_i |r_i| \text{ and } x^\dagger Q(\widehat{G})x \geq \lambda_{\min}(Q(\widehat{G})). \quad (6)$$

In order to show that $L_-(G)$ is positive semi-definite, for any non-zero unit vector $x \in \mathbb{C}^n$, we must have $x^\dagger L_-(G)x = x^\dagger R(G)x + x^\dagger Q(\widehat{G})x \geq 0$. If the given condition is satisfied, the proof follows from (6).

Given a weighted digraph, the lemma declares that if the maximum of the modulus of weights of the loops do not exceed the minimum signless Laplacian eigenvalue of the graph without considering the loops, then the signed Laplacian corresponding to the given graph will be positive semi-definite.

Example 2.10 1. Consider the graph given in Figure 1. The signed Laplacian matrix with negative loops associated with G is given by

$$\begin{aligned} L_-(G) &= \begin{bmatrix} 1 & 1 & 1 & -1 \\ 1 & 1 & 1 & -1 \\ 1 & 1 & 1 & -1 \\ -1 & -1 & -1 & 1 \end{bmatrix} \\ &= \begin{bmatrix} -2 & 0 & 0 & 0 \\ 0 & -2 & 0 & 0 \\ 0 & 0 & -2 & 0 \\ 0 & 0 & 0 & -2 \end{bmatrix} + \begin{bmatrix} 3 & 1 & 1 & -1 \\ 1 & 3 & 1 & -1 \\ 1 & 1 & 3 & -1 \\ -1 & -1 & -1 & 3 \end{bmatrix} \\ &= R(G) + Q(\widehat{G}). \end{aligned}$$

Note that, eigenvalues of $Q(\widehat{G})$ are 2, 2, 2, 6.




Figure 1: Graph with negative weighted loops

2. Consider the weighted digraph given in Figure 2 The eigenvalues of $L_-(G)$ are given by




Figure 2: Graph with negative weighted loops

0.0679, 1.8000, 3.5321.

Proposition 2.11 *Let G be a weighted digraph with loops having nonpositive weights. If the subgraph \widehat{G} obtained from G by removing the loops has signless Laplacian eigenvalue zero then $L_-(G)$ is not positive semi-definite.*

Proof: Let x be the unit eigenvector corresponding to the zero Laplacian eigenvalue of \widehat{G} . Then $x^\dagger L_-(G)x = x^\dagger R(G)x < 0$ since R contains at least one loop with negative weight.

Theorem 2.12 *Let G be a connected weighted digraph with nonpositive loop weights and n number of vertices. Let λ be a signless Laplacian eigenvalue of \widehat{G} of algebraic multiplicity k . Then the number of zero eigenvalues of $L_-(G)$ is k if $|r_i| = \lambda$ for all $i \in V(G)$.*

Proof: Consider $L_-(G) = R(G) + Q(\widehat{G})$. We know that 0 is an eigenvalue of $L_-(G)$ if and only if there exists a unit vector $x \in \mathbb{C}^n$ such that $x^\dagger L_-(G)x = 0$, which implies $x^\dagger R(G)x = x^\dagger Q(\widehat{G})x$. Assume that λ is an eigenvalue of $Q(\widehat{G})$ with algebraic multiplicity k . Since $Q(\widehat{G})$ is Hermitian, there exist unit orthogonal vectors x_1, x_2, \dots, x_k such that $x_i^\dagger Q(\widehat{G})x_i = \lambda$, $i = 1, \dots, k$. Set $R(G) = -\lambda I_n$, where I_n is the identity matrix of order n . Then $x_i^\dagger L_-(G)x_i = 0$ for all i . Since x_i s are orthonormal vectors, algebraic multiplicity of eigenvalue 0 of $L_-(G)$ is k .

Remark 2.13 *Note that, a connected weighted digraph with all negative weighted loops can have zero signed Laplacian eigenvalues of multiplicity more than one. In particular, consider any connected weighted complete digraph G where all the loops are present having weights equal to one of the eigenvalues of $Q(\widehat{G})$. For example, consider Figure 1.*

Now we consider weighted digraphs with loops having both positive and negative weights. Then we have the following lemma.

Lemma 2.14 *Let G be a weighted digraph with loops having at least one negative and at least one positive weighted loops. Assume that G has n vertices, k are having positive loop weights, l are having negative loop weights such that $k + l \leq n$. Let r_1^+, \dots, r_k^+ be the positive weights and r_{k+1}^-, \dots, r_l^- the negative weights such that $r_j^\pm = r_j - d_j$, for some $r_j \in \mathbb{R}$,*

$j = 1, \dots, k, k+1, \dots, l, \dots, n$ where d_j is the degree of j th vertex of \widehat{G} . Then $L_{\pm}(G)$ is positive semi-definite if and only if

$$\text{diag}\{r_1, r_2, \dots, r_n\} + A(\widehat{G})$$

is positive semi-definite.

Proof: Without loss of generality, assume that the first k vertices are having loops with positive weights, second l vertices are having loops with negative weights, and the remaining (if any) vertices have no loops. For any nonzero $x \in \mathbb{C}^n$,

$$\begin{aligned} x^\dagger L_{\pm}(G)x &= x^\dagger \begin{bmatrix} R_+ & 0 & 0 \\ 0 & R_- & 0 \\ 0 & 0 & R_0 \end{bmatrix} x + x^\dagger Q(\widehat{G})x \\ &= x^\dagger (\text{diag}\{r_1, r_2, \dots, r_n\} + A(\widehat{G}))x \end{aligned}$$

where $R_+ = \text{diag}\{r_1^+, \dots, r_k^+\}$, $R_- = \text{diag}\{r_1^-, \dots, r_l^-\}$ and $R_0 = \text{diag}\{r_{l+1}, \dots, r_n\}$. Hence, the result follows.

Theorem 2.15 *Let G be a weighted digraph with loops having both positive and negative weighted loops. The number of zero eigenvalues of $L_{\pm}(G)$ is equal to the number of zero eigenvalues of $\text{diag}\{r_1, r_2, \dots, r_n\} + A(\widehat{G})$ where $r_j^{\pm} = r_j - d_j, j = 1, \dots, n$ and r_j^{\pm} is the weight of the loop at j th vertex.*

Proof: The proof follows by Lemma 2.14.

Example 2.16 *Consider the weighted digraph in Figure 3. The signed Laplacian is*




Figure 3: Graph G with negative and positive weighted loops

$$L_{\pm}(G) = \begin{bmatrix} 1 & 0 & 0 & -2 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ -2 & 0 & 0 & 4 \end{bmatrix} = \begin{bmatrix} -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 2 \end{bmatrix} + \begin{bmatrix} 2 & 0 & 0 & -2 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ -2 & 0 & 0 & 2 \end{bmatrix}.$$

3 Graph structure for pure and mixed states

The approach of identifying the density matrix representations of quantum states by density matrices of unweighted undirected graphs was introduced in [5] and extended in [6] for weighted graphs. However, the recent development of signless Laplacian matrix associated with a graph has not been gainfully used in both [5] and [6] to define density matrix associated with the graph. Thus, the results in [6] could not capture interesting connections between the properties of density matrices defined by a graph and topology of the graph. In this section, we use the Laplacian matrices defined in the last section to define density matrix associated with a weighted digraph. Further, we describe how the topological structure of the graph dictates whether the corresponding density matrices represent pure or mixed states.

3.1 Pure States and Mixed States

Recall that density matrix representation of a quantum state is a Hermitian positive semi-definite matrix with unit trace. The density matrix σ corresponding to a state is called pure if $\text{Tr}(\sigma^2) = 1$ and mixed if $\text{Tr}(\sigma^2) < 1$. A quantum state, in general, can also be represented as

$$\sigma = \sum_i p_i |\psi_i\rangle\langle\psi_i|, \quad (7)$$

where $0 \neq |\psi_i\rangle \in \mathbb{C}^2$ with norm one and $\sum_i p_i = 1, 0 \leq p_i \leq 1$. Thus σ is a convex combination of rank one matrices, in particular, rank one projections. If σ is just a projection with rank one then σ is a pure state, otherwise, a mixed state.

We define density matrices associated with a weighted digraph. We denote an weighted complete bipartite digraph without loops of order n by K_n .

Definition 3.1 *The density matrix σ_G associated with an weighted digraph G are given by*

$$\sigma_G := \frac{1}{\text{Tr}(K(G))} K(G) \quad (8)$$

where

- $K(G) = L(G)$ when G is without a loop
- $K(G) = Q(G)$ when either G is without a loop or having loops with nonnegative weights
- $K(G) = L_-(G)$ when G is with loops having nonpositive weights and $L_-(G)$ is positive semi-definite
- $K(G) = L_\pm(G)$ when G contains loops with positive weights, negative weights and $L_\pm(G)$ is positive semi-definite.

Theorem 3.2 *The density matrix defined by Laplacian or signless Laplacian matrix of a weighted digraph G without loops has rank one if and only if the graph is K_2 or $\widehat{K}_2 := K_2 \sqcup v_1 \sqcup v_2 \sqcup \dots \sqcup v_{n-2}$, where v_1, v_2, \dots, v_{n-2} are isolated vertices.*

Proof: Assume that σ_G has rank one and G contains n vertices. Then σ_G has eigenvalue 1 with multiplicity one (since trace of $\sigma_G = 1$) and 0 is an eigenvalue of multiplicity $n - 1$. If $n = 2$ then obviously $G = K_2$. If $n \neq 2$ then by Corollary 2.7, G contains $n - 1$ connected components. Thus $G = \widehat{K}_2$.

Conversely, suppose $G = K_2$ or \widehat{K}_2 . Then the eigenvalues of σ_G are 0 with multiplicity $n - 1$ for $G = \widehat{K}_2$ and multiplicity 1 for $G = K_2$, and 1 with multiplicity one. Hence the result follows.

Remark 3.3 *We mention that the same result has been obtained in [5] for unweighted undirected graphs.*

Corollary 3.4 *Let G be a weighted digraph without loops isomorphic to K_2 or \widehat{K}_2 . Then σ_G constructed by $L(G)$ or $Q(G)$ represents a pure state.*

Proof: The density matrix σ_G has a simple eigenvalue 1 and other eigenvalues are zeros. Since trace of any matrix is sum of the eigenvalues of the matrix, we have $\text{Tr}(\sigma_G) = 1$ and $\text{Tr}(\sigma_G^2) = 1$. Thus the result follows.

Corollary 3.5 *Let G be a weighted digraph without loops of order n that is not isomorphic to K_2 and \widehat{K}_2 . Then σ_G constructed by $L(G)$ or $Q(G)$ represents a mixed state.*

Proof: Let the eigenvalues of $\sigma(G)$ be $\lambda_1 \leq \lambda_2 \leq \dots \leq \lambda_n$. By the definition of σ_G we have $\text{Tr}(\sigma_G) = 1$ as $\sum_{i=1}^n \frac{\lambda_i}{d(G)} = 1$ where $d(G) = \sum_{i=1}^n \lambda_i$. Then the eigenvalues of σ_G^2 are $\frac{\lambda_1^2}{d(G)^2}, \frac{\lambda_2^2}{d(G)^2}, \dots, \frac{\lambda_n^2}{d(G)^2}$. Thus

$$\text{Tr}(\sigma_G^2) = \frac{\sum_{i=1}^n \lambda_i^2}{d(G)^2} = \frac{d(G)^2 - 2 \sum_{i \neq j, i, j=1}^n \lambda_i \lambda_j}{d(G)^2} < 1.$$

Hence G represents a mixed state.

Example 3.6 1. Consider the graph in Figure 4 which represents a pure state.




Figure 4: Pure state given by K_2 along with 4 isolated nodes

2. Consider the graph in Figure 5 which represents a mixed state.




Figure 5: Mixed state with 6 vertices

The digraph O_1 denotes a digraph with one vertex and the vertex contains a directed weighted loop.

Theorem 3.7 The density matrix of order n defined by signless Laplacian matrix associated with a weighted digraph G with loops having nonnegative weights has rank one if and only if the graph is $\hat{O}_1 := O_1 \sqcup v_1 \sqcup v_2 \sqcup \dots \sqcup v_{n-1}$, where v_1, v_2, \dots, v_{n-1} are isolated vertices without loops.

Proof: Assume that the density matrix σ_G of G constructed by the signless Laplacian of the graph and G contains n vertices with at least one vertex contains a directed loop. Obviously, the matrix σ_G has rank 1 if and only if any submatrix of σ_G of order ≥ 2 is singular. Without loss of generality assume that the first vertex v_1 is attached with a directed loop. Then the following cases arise.

Case-I: The vertex v_1 is linked by a directed edge with another vertex say the second vertex v_2 . In this case, if we consider the 2×2 submatrix of σ_G constructed by the intersection of the 1st row, 2nd row, 1st column and 2nd column of σ_G then this is a matrix with nonzero determinant. Hence rank of σ_G is at least 2.

Case-II: Any two vertices without loops say v_i and v_j are linked by a directed edge. In this case, if we consider the 2×2 submatrix of σ_G constructed by the intersection of the 1st row, i th row, 1st column and i th column of σ_G then this is a matrix with nonzero determinant. Hence rank of σ_G is at least 2.

Case-III: All the vertices are isolated without loops except two vertices say the 1st and the 2nd vertex. Then, if we consider the 2×2 submatrix of σ_G constructed by the intersection of the 1st row, 2nd row, 1st column and 2nd column of σ_G then this is a matrix with nonzero determinant. Hence rank of σ_G is at least 2.

Case-IV: All the vertices are isolated without loops except the 1st vertex. Then the density matrix σ_G is of rank 1.

Hence the desired result follows.

Corollary 3.8 Let G be a weighted digraph with loops having nonnegative weights and is isomorphic to \hat{O}_1 . Then σ_G defined by the signless Laplacian of G represents a pure state.

Corollary 3.9 *Let G be a weighted digraph with loops having nonnegative weights of order n that is not isomorphic to \widehat{O}_1 . Then σ_G defined by the signless Laplacian of G represents a mixed state.*

Example 3.10 1. *Consider the graph in Figure 6 which represents a pure state.*



Figure 6: Pure state given by O_1 along with a single loop

2. *Consider the graph in Figure 7 which represents a mixed state.*




Figure 7: Mixed state with 6 vertices

Remark 3.11 *Realization of 1-qubit by weighted digraph: Consider the graph $G = K_2$ for a pure state with edge weight $w \in \mathbb{S}_1^+$. Then the corresponding density matrix with respect to the Laplacian matrix is given by*

$$\sigma_G = \frac{1}{2}L(G) = \frac{1}{2} \begin{bmatrix} 1 & -w \\ -\bar{w} & 1 \end{bmatrix},$$

where $w = e^{i\phi}$, $0 \leq \phi \leq 2\pi$. The eigenvalues of σ_G are 0 and 1 corresponding to eigenvectors $|\psi_1\rangle = \frac{1}{\sqrt{2|z_1|}} \begin{bmatrix} z_1 \\ \bar{w}z_1 \end{bmatrix}$ and $|\psi_2\rangle = \frac{1}{\sqrt{2|z_2|}} \begin{bmatrix} z_2 \\ -\bar{w}z_2 \end{bmatrix}$ respectively, where $0 \neq z_1, z_2 \in \mathbb{C}$. Thus the pure state is given by $\sigma = |\psi_2\rangle\langle\psi_2|$. Setting $z_2 = re^{i\theta}$, $|z_2| = r > 0$, $0 \leq \theta \leq 2\pi$, the vector representation of the pure state is given by

$$\begin{aligned} |\psi\rangle &= e^{i\theta} \left(\frac{1}{\sqrt{2}}|0\rangle - \frac{1}{\sqrt{2}}e^{-i\phi}|1\rangle \right), \\ &\equiv \frac{1}{\sqrt{2}}|0\rangle - \frac{1}{\sqrt{2}}e^{-i\phi}|1\rangle \end{aligned}$$

where $|0\rangle = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$ and $|1\rangle = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$.

Further, the density matrix with respect to the signless Laplacian matrix is given by

$$\sigma_G = \frac{1}{2}Q(G) = \frac{1}{2} \begin{bmatrix} 1 & w \\ \bar{w} & 1 \end{bmatrix}.$$

Following a similar approach, as above, the corresponding vector representation of the pure state is given by

$$|\psi\rangle \equiv \frac{1}{\sqrt{2}}|0\rangle + \frac{1}{\sqrt{2}}e^{-i\phi}|1\rangle, 0 \leq \phi \leq 2\pi.$$

Now we consider weighted graphs with loops having nonpositive weights. We denote S_n , $n \geq 2$, a star graph with n vertices.

Theorem 3.12 *Consider a weighted digraph G consisting of a weighted digraph \widehat{G} without loops having n number of vertices and loops at each vertex of \widehat{G} with equal weights $-\lambda$ where λ is a signless Laplacian eigenvalue of \widehat{G} with multiplicity $n - 1$. Then $\sigma(G) = \frac{1}{\text{Tr}(L_-(G))}L_-(G)$ represents a pure state.*

Proof: By Lemma 2.9, $\sigma(G)$ is positive semi-definite. However, $n - 1$ number of eigenvalues of $L_-(G)$ are zero since λ is an eigenvalue of $Q(\widehat{G})$ of algebraic multiplicity $n - 1$. Therefore, rank of $\sigma(G)$ is one. Hence the result follows.

Corollary 3.13 Consider a weighted digraph G consists of a weighted digraph \widehat{G} without loops having n number of vertices and loops at each vertex of \widehat{G} with equal weights $-\lambda$ where λ is a signless Laplacian eigenvalue of \widehat{G} with multiplicity $k < n - 1$. Then $\sigma(G) = \frac{1}{\text{Tr}(L_-(G))}L_-(G)$ represents a mixed state.

Example 3.14 1. Consider $G = K_n$ along with loops at each vertex of equal weights $-n/(n - 1)$. Then $\sigma(G) = \frac{1}{\text{Tr}(L_-(G))}L_-(G)$ represents a pure state. For instance, consider $n = 3$ in Figure 8.




Figure 8: Pure state given by K_3 along with negative weighted loops

2. Consider $G = S_n$ along with loops at each vertex of equal weights -1 . Then $\sigma(G) = \frac{1}{\text{Tr}(L_-(G))}L_-(G)$ represents a mixed state. For example, consider $n = 4$ in Figure 9.




Figure 9: Mixed state given by S_4 along with negative weighted loops

Now we consider graphs with both positive and negative weighted loops.

Theorem 3.15 Let G be a weighted digraph with loops having at least one negative and at least one positive weighted loops. Assume that G has n vertices, k are having positive loop weights, l are having negative loop weights such that $k + l \leq n$. Let r_1^+, \dots, r_k^+ be the positive weights and r_{k+1}^-, \dots, r_l^- the negative weights such that $r_j^\pm = r_j - d_j, j = 1, \dots, k, k + 1, \dots, l, \dots, n$ where d_j is the degree of j th vertex of \widehat{G} . Then G represents a pure state if and only if

$$\text{diag}\{r_1, r_2, \dots, r_n\} + A(\widehat{G})$$

has rank one.

Proof: The proof follows from the construction of $L_\pm(G)$ and Lemma 2.14.

However, we can construct a class of pure states by using the construction mentioned in the following corollary.

Corollary 3.16 Consider a digraph \widehat{G} without loops having n vertices that represents a pure state obtained by signless Laplacian matrix $Q(\widehat{G})$, that is, only two vertices of G , say i th and j th of $\widehat{G} = (V, E)$ are linked having edge weight $w_{ij} \in \mathbb{C}$, rest of the vertices are isolated.

Define $r_i^+ = r_i^2 - |w_{ij}|$ and $r_j^- = r_j^2 - |w_{ij}|$ where $r_i, r_j \in \mathbb{R}_+$ such that $r_i^2 + r_j^2 = 1$ and $r_i r_j = |w_{ij}|$. Then the graph G constructed by \widehat{G} along with loops introduced at the i th and j th vertices having weights r_i^+ and r_j^- respectively, provides a pure state defined by $L_{\pm}(G)$.

Proof: Note that, all the entries of $L_{\pm}(G) = (l_{pq})$ are given by

$$(L_{\pm}(G))_{pq} = \begin{cases} r_i^2, & \text{if } p = q = i \\ r_j^2, & \text{if } p = q = j \\ w_{ij}, & \text{if } p = j, q = i \\ \overline{w_{ij}}, & \text{if } p = i, q = j \\ 0 & \text{otherwise.} \end{cases}$$

Obviously, $L_{\pm}(G)$ is Hermitian, positive semi-definite and $\text{Tr}(L_{\pm}(G)) = 1$. Further, rank of $L_{\pm}(G) = 1$. Therefore, G represents a pure state.

4 Graph structure of entangled pure states

In this section, we provide weighted digraphs whose density matrices represent entangled pure states. Because of the potential applications offered by pure entangled states, they are of immense importance in quantum information and computation. This forms the basis for studying the properties of such states from a graph theoretic approach.

1. *Bell States:* For two-qubit systems, Bell states [1] are maximally entangled states represented as

$$\begin{aligned} |\phi\rangle_{12}^{\pm} &= \frac{1}{\sqrt{2}} [|00\rangle_{12} \pm |11\rangle_{12}], \\ |\psi\rangle_{12}^{\pm} &= \frac{1}{\sqrt{2}} [|01\rangle_{12} \pm |10\rangle_{12}]. \end{aligned} \quad (9)$$

For example, consider the graphs with four vertices in Figure 10. The density matrices



Figure 10: G_1 and G_2

are given by $\sigma(G_i) = \frac{K(G_i)}{\text{Tr}(K(G_i))}$; $K(G_i) \in \{L(G_i), Q(G_i)\}$ where $\sigma(G_1) = |\phi\rangle_{12}^+ \langle\phi|_{12}^+$ and $\sigma(G_2) = |\psi\rangle_{12}^+ \langle\psi|_{12}^+$. In order to produce the Bell states of the form $\frac{1}{\sqrt{2}} [|00\rangle + e^{i\delta} |11\rangle]$ and $\frac{1}{\sqrt{2}} [|01\rangle + e^{i\delta} |10\rangle]$ using G_1 and G_2 , one has to replace the edge weights by a factor $e^{i\delta}$ and edge will be unidirectional.

2. *General 2-qubit and 3-qubit entangled states:* Consider the graph in Figure 11 for the




Figure 11: Graph G

general two-qubit state $|\Psi\rangle = a |00\rangle + b |11\rangle$, where $a, b \in \mathbb{C} \setminus \{0\}$ and $|a|^2 + |b|^2 = 1$. The density matrix associated with G is given by $\sigma(G) = \frac{L_{\pm}(G)}{\text{Tr}(L_{\pm}(G))} = |\Psi\rangle \langle\Psi|$. The graph for a general 3-qubit state, $|\Phi\rangle = a |000\rangle + b |111\rangle$, will follow similarly by considering 8 vertices where only the first and the last vertices will be linked.

3. *Three-qubit GHZ and W States:* Three qubit states can be separated into two inequivalent classes, namely GHZ class and W class [15, 16]. These classes have distinct properties and cannot be converted into one another by performing Stochastic Local Operations and Classical Communication (SLOCC). We have already shown that the graph of a general three-qubit GHZ state will be similar to Fig. (11). For specific cases of $a = b = \frac{1}{\sqrt{2}}$, the eight orthogonal GHZ states are,

$$\begin{aligned}
|\psi\rangle_{123}^{(1),(2)} &= \frac{1}{\sqrt{2}} [|000\rangle \pm |111\rangle], \\
|\psi\rangle_{123}^{(3),(4)} &= \frac{1}{\sqrt{2}} [|001\rangle \pm |110\rangle], \\
|\psi\rangle_{123}^{(5),(6)} &= \frac{1}{\sqrt{2}} [|010\rangle \pm |101\rangle], \\
|\psi\rangle_{123}^{(7),(8)} &= \frac{1}{\sqrt{2}} [|011\rangle \pm |100\rangle],
\end{aligned} \tag{10}$$

The graphs corresponding to Eq. (12) are given in Figures 12,13, 14, 15.




Figure 12: G_1



Figure 13: G_2




Figure 14: G_3




Figure 15: G_4

and the density matrices associated with G_1, G_2, G_3, G_4 are given $\sigma(G_i) = \frac{K(G_i)}{\text{Tr}(K(G_i))}$ where $K(G_i) \in \{L(G_i), Q(G_i)\}$, and $\sigma(G_1) = |\psi\rangle_{123}^{(1),(2)} \langle\psi|_{123}^{(1),(2)}$, $\sigma(G_2) = |\psi\rangle_{123}^{(3),(4)} \langle\psi|_{123}^{(3),(4)}$, $\sigma(G_3) = |\psi\rangle_{123}^{(5),(6)} \langle\psi|_{123}^{(5),(6)}$, $\sigma(G_4) = |\psi\rangle_{123}^{(7),(8)} \langle\psi|_{123}^{(7),(8)}$.

The general three-qubit W state is given as $|\psi\rangle_{123}^W = a|001\rangle + b|010\rangle + c|100\rangle$ where $|a|^2 + |b|^2 + |c|^2 = 1$. The graph representation of $|\psi\rangle_{123}^W$ is given in Figure 16.

The density matrix for the W class of states can be expressed as $\sigma(G) = \frac{L-(G)}{\text{Tr}(L-(G))}$. For a specific case where $a = b = c = \frac{1}{\sqrt{3}}$, Figure 17 represents the graphical representation for a standard W state. It is evident that the graphs for GHZ and W classes are completely distinct from each other. Therefore, using our approach one can easily identify whether a given three-qubit state belongs to a GHZ class or W class.




Figure 16: G




Figure 17: G

Similar to $G_i, i = 1, \dots, 4$ graphs with 8 vertices, graphs with 16 vertices can be produced which will provide the GHZ states with 4-qubits. By similar graphs, we mean graphs with 16 nodes, one edge which connects the i th and $(17 - i)$ th vertices, $i = 1, \dots, 8$ having edge weight 1. Similarly one can also obtain the graph for a four qubit W state.

4. *Cluster and Chi states:* The four qubit cluster [17] and Chi [18] states are given by $|\psi\rangle_{1234} = \frac{1}{2}(|0000\rangle + |0101\rangle + |1010\rangle - |1111\rangle)$ and $|\phi\rangle_{1234} = \frac{1}{2}(|0000\rangle + |0101\rangle + |1011\rangle - |1110\rangle)$, respectively. The corresponding graphs for these two states are given in Figures 18 and 19, respectively. The two states are different as evident from the edge weights.



Figure 18: G




Figure 19: G

Similarly, the density matrices corresponding to Cluster and Chi states are represented by $\sigma(G_i) = \frac{L_-(G_i)}{\text{Tr}(L_-(G_i))}$ where $\sigma(G_1) = |\psi\rangle_{1234} \langle\psi|_{1234}$, and $\sigma(G_2) = |\phi\rangle_{1234} \langle\phi|_{1234}$.

5. *Brown State:* Consider the graph G with 32 vertices given in Figure (20). The isolated vertices are not shown in the graph and the weights of the edges and loops are as given

below.

$$w_{ij} = \begin{cases} -1, & i = 1, j = 4, 15 \\ 1, & i = 1, j = 14, 21, 24, 26, 27 \\ -1, & i = 4, j = 14, 21, 24, 26, 27 \\ 1, & i = 4, j = 15 \\ -1, & i = 14, j = 15 \\ 1, & i = 14, j = 21, 24, 26, 27 \\ -1, & i = 15, j = 21, 24, 26, 27 \\ 1, & i = 21, j = 24, 26, 27 \\ 1, & i = 24, j = 26, 27 \\ 1, & i = 26, j = 27 \\ -6, & i = j, i = 1, 4, 14, 15, 21, 24, 26, 27. \end{cases} \quad (11)$$

The graph in Figure (20) represents a five qubit Brown state [19], namely

$$\begin{aligned} |\psi\rangle^{12345} &= \frac{1}{2\sqrt{2}} [|00000\rangle - |00011\rangle + |01101\rangle - |01110\rangle + |10100\rangle + |10111\rangle + |11001\rangle \\ &+ |11010\rangle] \end{aligned} \quad (12)$$

In comparison to other non-equivalent classes of five-qubit entangled states, Brown states are said to be more entangled. The reason is evident from the property that all the bipartitions of Brown states are maximally mixed which is not the case with GHZ, Cluster or Chi type of states.

The density matrix associated with the graph for Brown state is

$$\sigma(G) = \frac{1}{\text{Tr}(L_-(G))} L_-(G) = |\psi\rangle_{12345} \langle\psi|_{12345}.$$




Figure 20: G

6. *5-qubit Chi state*: The five-qubit Chi state [20] can be represented as

$$\begin{aligned} |\phi\rangle_{12345} &= \frac{1}{2} [|00000\rangle + |00111\rangle + |01010\rangle - |01101\rangle - |10011\rangle + |10100\rangle + |11001\rangle \\ &+ |11110\rangle]. \end{aligned} \quad (13)$$

The graph and weights of edges for the five-qubit Chi state are given by Figure (21) and Eq. (14), respectively.




Figure 21: G

$$w_{ij} = \begin{cases} -1, & i = 1, j = 14, 20 \\ 1, & i = 1, j = 8, 11, 21, 26, 31 \\ -1, & i = 8, j = 14, 20 \\ 1, & i = 8, j = 11, 21, 26, 31 \\ -1, & i = 11, j = 14, 20 \\ 1, & i = 11, j = 21, 26, 31 \\ -1, & i = 14, j = 21, 26, 31 \\ 1, & i = 14, j = 20 \\ -1, & i = 20, j = 21, 26, 31 \\ 1, & i = 26, j = 31 \\ -6, & i = j, i = 1, 4, 14, 15, 21, 24, 26, 27. \end{cases} \quad (14)$$

The density matrix, therefore, can be given as $\sigma(G) = \frac{1}{\text{Tr}(L_-(G))} L_-(G) = |\phi\rangle_{12345} \langle\phi|_{12345}$.

Although the difference between Brown and Chi states can be characterized from the edge weights, for a meaningful classification of such states using a graph theoretical approach, one needs to quantify a graph theoretic measure for entanglement. Such a measure will classify quantum states in different classes and provide deeper physical insight into the complex nature of multiqubit entanglement.

Remark 4.1 *Observe that in the graph representation of entangled pure states mentioned above, all the existing weighted edges are clustered in a completely connected subgraph of the original graph. Further, the weight of the loops attached at each of the vertices of the subgraph is $-(m-2)$ where m is the number of vertices involved in the complete subgraph.*

5 Conclusion

We define combinatorial, signless and signed Laplacian matrices associated with a weighted digraph having complex edge weights with or without loops. We determine the connection between the existence of zero Laplacian eigenvalues of a weighted digraph and the topological structure of the graph. Using these Laplacian matrices, we define density matrices corresponding to a weighted digraph. We have classified graphs which represent pure and mixed density matrices of quantum states by using the topological structure of the graphs. This work initiates a number of directions to the combinatoric visualization of quantum mechanical phenomena. Some of them are listed below.

1. A state is called separable if the density matrix,

$$\rho = \sum_i p_i \rho_i^{(A)} \otimes \rho_i^{(B)}.$$

Here, $\rho_i^{(A)}$ and $\rho_i^{(B)}$ denotes density matrix of subsystems A and B . We denote tensor product of matrices by \otimes . The state is entangled otherwise. We have deduced the graphs for several well-known entangled pure states. We have demonstrated that the three-qubit entangled systems can be classified into GHZ and W class using a graph theoretic approach. A criteria of separability states represented by the Laplacian of simple graphs has been developed in [7]. A combinatorial operation has also been introduced for density matrices defined by Laplacian matrices associated with simple graphs in [9] that act as an entanglement generator for mixed states arising from partially symmetric graphs. These works introduce new results for the separability of density matrices corresponding to weighted digraphs.

2. In order to develop further insight into the entanglement properties of multiqubit systems, it would be interesting to define a graph theoretic measure for quantification and classification of entanglement in such systems.
3. Recently, local unitary transformations on a density matrix obtained by signless Laplacian matrix associated with a simple graph has been established as a combinatorial operation which is known as switching of a graph in [8]. This work sheds further light to the problem of unitary equivalence and state classification for the states related to weighted digraphs.

This work is, we hope, a contribution towards a new direction in the field of quantum information.

References

- [1] A. Einstein, B. Podolsky, and N. Rosen, *Can Quantum-Mechanical Description of Physical Reality Be Considered Complete?*, Phys. Rev. **47**, 777-780 (1935); ; J. S. Bell, *On the Einstein-Podolsky-Rosen paradox*, Physics (Long Island City, N. Y.) **1**, 195-200 (1964).
- [2] W. K. Wootters, *Entanglement of formation of an arbitrary state of two qubits*, Phys. Rev. Lett. **80** (1998) 2245-2248.
- [3] A. Miyake, *Classification of multipartite entangled states by multidimensional determinants*, Phys. Rev. A **67** (2003) 012108: 1-10.
- [4] T. Sunada, *A discrete analogue of periodic magnetic Schrödinger operators*, *Geometry of the spectrum*, Contemp. Math., Amer. Math. Soc., Providence, RI (Seattle, WA, 1993), **173** (1994) 283-299.
- [5] S. L. Braunstein, S. Ghosh and S. Severini, *The Laplacian of a graph as a density matrix: a basic combinatorial approach to separability of mixed states*, Annals of Combinatorics **10** (2006) 291-317.
- [6] Hassan Ali, Saif M. and Pramod S. Joag. *A combinatorial approach to multipartite quantum systems: basic formulation*, Journal of Physics A: Mathematical and Theoretical **40.33** (2007): 10251.
- [7] Wu, Chai Wah, *Multipartite separability of Laplacian matrices of graphs*, Electronic journal of combinatorics **16.1** (2009): R61.
- [8] Dutta, Supriyo, Bibhas Adhikari and Subhashish Banerjee, *A graph theoretical approach to states and unitary operations*. Quantum Information Processing **15.5** (2016): 2193-2212.

- [9] Dutta, Supriyo, Bibhas Adhikari, Subhashish Banerjee and R. Srikanth, *Bipartite separability and non-local quantum operations on graphs*, arXiv preprint arXiv:1601.07704 (2016).
- [10] Reff, Nathan, *Spectral properties of complex unit gain graphs*, Linear Algebra and its Applications 436.9 (2012): 3165-3176.
- [11] R. B. Bapat, *Graphs and Matrices*, Hindustan Book Agency, New Delhi, India, Ist Edition (2011).
- [12] R. B. Bapat, D. Kalita and S. Pati, *On weighted directed graphs*, Linear Algebra and its Appl. 436.1 (2012): 99-111.
- [13] Cvetkovic Drago, Peter Rowlinson, and Slobodan K. Simic, *Signless Laplacians of finite graphs*, Linear algebra and its applications 423.1 (2007): 155-171.
- [14] Wu, Chai Wah, *Conditions for Separability in Generalized Laplacian Matrices and Diagonally Dominant Matrices as Density Matrices*, IBM Research Report RC23758(W0508-118)(October 18, 2005).
- [15] D. M. Greenberger, M. A. Horne., A. Shimony and A. Zeilinger, *Bell's theorem without inequalities*, A. J. Phys. **58**, 1131-1143 (1990);
- [16] W. Dur, G. Vidal and J. I. Cirac, Phys. Rev. A **62**, 062314-062314(12) (2000).
- [17] H.J. Briegel and R. Raussendorf, *Persistent Entanglement in arrays of Interacting Particles*, Phys. Rev. Lett. 86, 910-913 (2001).
- [18] Y. Yeo and W. K. Chua, *Teleportation and Dense Coding with Genuine Multipartite Entanglement*, Phys. Rev. Lett. 96, 060502(1)-060502(4) (2006).
- [19] I. D. K. Brown, S. Stepney, A. Sudbery, and S. L. Braunstein, *Searching for highly entangled multi-qubit states*, J. Phys. A 38, 1119-1131, (2005).
- [20] Z. X. Man, Y. J. Xia, and N. Ba An, *Genuine multiqubit entanglement and controlled teleportation*, Phys. Rev. A. 75, 05306(1)-05306(5) (2006)